


# Turbulent seismoacoustic signals from a hurricane landfall


Qing Ji<sup>1</sup> Ipshita Dey<sup>2,3</sup> Eric Dunham<sup>1</sup>

- 1 Department of Geophysics, Stanford University
- 2 Department of Earth System Science, Stanford University
- 3 National Centre for Atmospheric Science and Department of Meteorology, University of Reading, UK











#### Infrasound data for hurricane study




Rose Stiale Contain ways Infonitaring of the evaluation to when the supply ting quartifles (e.g., wind speed, turbulence analysis, ...)

#### As hurricane passes the station .....




#### As hurricane passes the station .....




#### Wavelet spectrograms of infrasound & seismic data

1000



Ji & Dunham (2024)


## High coherence indicates local quasi-static response




#### **Simplified illustration**

Sorrells (1971) theory


Pressure wave model




#### Interdisciplinary modeling



## LES of Hurricane Boundary Layer (HBL) on land



#### **Turbulent origin of signals**




1. Insights into Sorrells theory


2. Infrasound data for turbulence analysis

Ji et al. (in revision)

#### Convection velocity of a turbulent field

Turbulent boundary layer







Velocity is zero at the wall, but wall stress / pressure advects with  $U_c$  (convection velocity) "Shadow" of eddies at height where wind speed  $U = U_c \approx 0.6 U_0$ 

Sorrells theory: "Pressure wave speed"  $c \approx U_c$ 

Similar in Mars application, where  $c \approx 0.5 U_0$  work best for Sorrells results (Murdoch et al. 2017)

#### Turbulent spectra from wind tower and infrasound





#### **Streamwise velocity PSD**

$$S_{uu}(f) = \alpha_u \left(\frac{\varepsilon U}{2\pi}\right)^{2/3} f^{-5/3}$$


**Dissipation rate &** is a key parameter describing the turbulence statistic, and contributes to an important energy source for hurricanes. (Bister & Emanuel, 1998)

#### **Pressure PSD**

$$\frac{1}{\rho^2} S_{pp}(f) = \alpha_p \left(\frac{\varepsilon U_c}{2\pi}\right)^{4/3} f^{-7/3}$$

Kolmogorov (1941), Obukhov (1949), George et al. (1984) ...

## Continuous monitoring of hurricane landfall




Fluctuation:  $p_{rms} \propto u_{rms}^2$ 

Turbulent intensity:  $u_{rms} \propto U_{10 \text{ m}}$ 

Fit wind model:  $U_{10 \text{ m}} = U(r)$ 


ERA5 reanalysis has a larger radius of maximum wind.



 $O(0.1 \text{ m}^2/\text{s}^3)$  or O(0.1 W/kg) dissipation rate

Interpreted as  $\varepsilon$  for height  $\sim 100$  m where the wind speed equal to  $U_c$ 

Ji et al. (in revision)



# Turbulent dissipation rate during hurricane landfall

For **pressure** spectra:


Assume  $U_c \approx 1.8 \ U_{10 \text{ m}}$  from our LES result Directly use  $U_{10 \text{ m}} \approx 4.2 \sqrt{p_{\text{rms}}}$  in their units

Wind tower for 15 m high, while pressure imprints from ~100 m high (where  $U = U_c$ )

Magnitude of  $\varepsilon$  agrees with other studies on landfalling storms.

Scaling  $\varepsilon \propto U^3$  considered for dominance of shear production, but less certain for storms.





TA.142A, 9.6 km

Aug 29

Aug 28

Aug 30

Aug 31

ASOS (LA.MLU

Aug 26

Aug 26

Aug 27

Aug 28

Aug 29

Aug 30

TA.140A, 17.5 km

ERA5 (TA.140A)

ASOS (LA.BAD)

Aug 26

Aug 27

Aug 27 Aug 28


Aug 28


Aug 29

Aug 30

Aug 29 Aug 30 Aug 31 Sep 01

Aug 31





Lag time (s)

 $D_{pp}$  (Pa<sup>2</sup>)

TA.341A

## **Turbulent pressure**

Structure function

$$D_{pp}(\tau) = [p(t+\tau) - p(t)]^{2}$$
$$\frac{1}{\rho^{2}}D_{pp}(\tau) = C_{p} (\varepsilon U_{c})^{4/3} \tau^{4/3}$$

Deviation from the slope:

- Violation of locally isotropic turbulence assumption
- Interaction of different scales

George et al. (1984), Katul et al. (2025)

Surprise #1: Seismic stations record in-situ data of Hurricane Issac after landfall

Distinct seismic ground motion contributed by ocean and atmosphere

Surprise #2: Turbulence explains the seismoacoustic signatures (in the atm. band)

Interdisciplinary modeling framework to explain observations

Surprise #3: Potential of seismoacoustic stations for environmental monitoring

Station networks with years of continuous data, especially infrasound

